



# 9<sup>th</sup> International Congress of Dipterology

**Abstracts Volume**

**25–30 November 2018  
Windhoek  
Namibia**

## **Organising Committee:**

Ashley H. Kirk-Spriggs (Chair)  
Burgert S. Muller  
Mary K. Kirk-Spriggs  
Gillian Maggs-Kölling  
Kenneth Uiseb  
Seth Eiseb  
Michael Osae  
Sunday Ekesi  
Candice-Lee Lyons

## **Edited by:**

Ashley H. Kirk-Spriggs  
Burgert S. Muller

# 9<sup>th</sup> International Congress of Dipterology

25–30 November 2018  
Windhoek, Namibia

## Abstract Volume

Edited by:

Ashley H. Kirk-Spriggs & Burgert S. Muller



Namibian Ministry of  
Environment and Tourism



NAMIBIA



WISSENSCHAFTLICHE GESELLSCHAFT

## Organising Committee

Ashley H. Kirk-Spriggs (Chair)  
Burgert S. Muller  
Mary K. Kirk-Spriggs  
Gillian Maggs-Kölling  
Kenneth Uiseb  
Seth Eiseb  
Michael Osae  
Sunday Ekesi  
Candice-Lee Lyons

Published by the International Congresses of Dipterology, © 2018.

Printed by John Meinert Printers, Windhoek, Namibia.

ISBN: 978-1-86847-181-2

**Suggested citation:** Adams, Z.J. & Pont, A.C. 2018. In celebration of Roger Ward Crosskey (1930–2017) – a life well spent. *In*: Kirk-Spriggs, A.H. & Muller, B.S., eds, *Abstracts volume. 9<sup>th</sup> International Congress of Dipterology, 25–30 November 2018, Windhoek, Namibia*. International Congresses of Dipterology, Windhoek, p. 2. [Abstract].

**Front cover image:** Tray of micro-pinned flies from the Democratic Republic of Congo (photograph © K. Pannecoucke).

**Cover design:** Craig Barlow (previously National Museum, Bloemfontein).

Disclaimer: Following recommendations of the various nomenclatorial codes, this volume is not issued for the purposes of the public and scientific record, or for the purposes of taxonomic nomenclature, and as such, is not published in the meaning of the various codes. Thus, any nomenclatural act contained herein (e.g., new combinations, new names, etc.), does not enter biological nomenclature or pre-empt publication in another work.

## A pilot study to delimit *tsetse* target populations in Zimbabwe

Gerald Chikowore<sup>1\*</sup>, Ahmadou H. Dicko<sup>2</sup>, Peter Chinwada<sup>3</sup>, Mosesimba<sup>3</sup>, William Shereni<sup>1</sup>, François Roger<sup>2</sup>, Jérémy Bouyer<sup>4</sup> & Laure Guerrini<sup>2</sup>

<sup>1</sup>Tsetse Control Division, Department of Livestock and Veterinary Services, Ministry of Lands, Agriculture and Rural Resettlement, P.O. Box CY52, Causeway, Harare, Zimbabwe. \*gkchikore@gmail.com

<sup>2</sup>Centre de Coopération Internationale en Recherche Agronomique pour le Développement, Montpellier, France.

<sup>3</sup>University of Zimbabwe, Harare, Zimbabwe.

<sup>4</sup>International Atomic Energy Agency, Vienna, Austria.

**Keywords:** Probability model, species distribution, *tsetse*, Zimbabwe.

*Tsetse* (Glossinidae) are cyclical vectors of human and animal trypanosomoses currently targeted by the African Union, under the Pan African Tsetse and Trypanosomiasis Eradication Campaign (PATTEC). Elaborate plans to guide intervention are a requirement if effective control of *tsetse* is to be achieved. A model to aid the planning of intervention programmes and assist a fuller understanding of *tsetse* distribution was applied in a pilot study covering 400 km<sup>2</sup> in Masoka, Zimbabwe and targeted two savanna species, *Glossina morsitans morsitans* Westwood and *Glossina pallidipes* Austen. Current data were used to study habitat suitability of both species, based on climatic and environmental data derived from MODIS and SPOT 5 satellite images. Factors influencing distribution were explored, using an Ecological Niche Factor Analysis (ENFA), whilst habitat suitability was predicted using a Maximum Entropy (MaxEnt) model at a spatial resolution of 250 m. Area Under the Curve (AUC), an indicator of model performance, was 0.89 for *G. m. morsitans* and 0.96 for *G. pallidipes*. The probability that flies were really absent from grid cells where they were not captured during the study was then calculated, based on a probability model using a risk threshold of 0.05. Apart from grid cells where *G. m. morsitans* and *G. pallidipes* were captured in the study area, there was a high probability of presence in additional grid cells adding up to 128 km<sup>2</sup> and 144 km<sup>2</sup> respectively. The modelling process promised to be useful in optimising the outputs of presence/absence surveys, allowing the definition of *tsetse* infested areas with improved accuracy.